

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1626GMS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 SEP 01 New pricing for the Save Answers for SciFinder Wizard within
STN Express with Discover!
NEWS 4 OCT 28 KOREAPAT now available on STN
NEWS 5 NOV 30 PHAR reloaded with additional data
NEWS 6 DEC 01 LISA now available on STN
NEWS 7 DEC 09 12 databases to be removed from STN on December 31, 2004
NEWS 8 DEC 15 MEDLINE update schedule for December 2004
NEWS 9 DEC 17 ELCOM reloaded; updating to resume; current-awareness
alerts (SDIs) affected
NEWS 10 DEC 17 COMPUAB reloaded; updating to resume; current-awareness
alerts (SDIs) affected
NEWS 11 DEC 17 SOLIDSTATE reloaded; updating to resume; current-awareness
alerts (SDIs) affected
NEWS 12 DEC 17 CERAB reloaded; updating to resume; current-awareness
alerts (SDIs) affected
NEWS 13 DEC 17 THREE NEW FIELDS ADDED TO IFIPAT/IFIUDB/IFICDB
NEWS 14 DEC 30 EPFULL: New patent full text database to be available on STN
NEWS 15 DEC 30 CAPLUS - PATENT COVERAGE EXPANDED
NEWS 16 JAN 03 No connect-hour charges in EPFULL during January and
February 2005
NEWS 17 FEB 25 CA/CAPLUS - Russian Agency for Patents and Trademarks
(ROSPATENT) added to list of core patent offices covered
NEWS 18 FEB 10 STN Patent Forums to be held in March 2005
NEWS 19 FEB 16 STN User Update to be held in conjunction with the 229th ACS
National Meeting on March 13, 2005
NEWS 20 FEB 28 PATDPAFULL - New display fields provide for legal status
data from INPADOC
NEWS 21 FEB 28 BABS - Current-awareness alerts (SDIs) available
NEWS 22 FEB 28 MEDLINE/LMEDLINE reloaded
NEWS 23 MAR 02 GBFULL: New full-text patent database on STN
NEWS 24 MAR 03 REGISTRY/ZREGISTRY - Sequence annotations enhanced
NEWS 25 MAR 03 MEDLINE file segment of TOXCENTER reloaded

NEWS EXPRESS JANUARY 10 CURRENT WINDOWS VERSION IS V7.01a, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 10 JANUARY 2005

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN

03/14/2005 10807710.trn

NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 14:16:29 ON 14 MAR 2005

=>

Uploading

THIS COMMAND NOT AVAILABLE IN THE CURRENT FILE

Do you want to switch to the Registry File?

Choice (Y/n):

Switching to the Registry File...

Some commands only work in certain files. For example, the EXPAND command can only be used to look at the index in a file which has an index. Enter "HELP COMMANDS" at an arrow prompt (=>) for a list of commands which can be used in this file.

=> FILE REGISTRY

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 14:16:38 ON 14 MAR 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 13 MAR 2005 HIGHEST RN 845467-46-1

DICTIONARY FILE UPDATES: 13 MAR 2005 HIGHEST RN 845467-46-1

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:

<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>


Uploading C:\Program Files\Stnexp\Queries\10807710.str

α -Toluenethiol, α -[1-(dimethylamino)ethyl]-, hydrochloride,
L(+)-threo-
(preparation of)

RN 942-48-3 CAPLUS

CN Benzenemethanethiol, α -[1-(dimethylamino)ethyl]-, [S-(R*,R*)]- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 942-49-4 CAPLUS

CN α -Toluenethiol, α -[1-(dimethylamino)ethyl]-, L-(-)-erythro-
(8CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 942-50-7 CAPLUS

CN Benzenemethanethiol, α -[1-(dimethylamino)ethyl]-, hydrochloride,
[S-(R*,R*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl

L13 ANSWER 17 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1964:447807 CAPLUS

DOCUMENT NUMBER: 61:47807

ORIGINAL REFERENCE NO.: 61:8284a-b



TITLE: Preparation of quaternary ammonium betaine salts

INVENTOR(S): Klass, Donald L.

PATENT ASSIGNEE(S): Pure Oil Co.

SOURCE: 4 pp.

DOCUMENT TYPE: Patent



chain nodes :

1 2 3 4 5 6 7 8 9

chain bonds :

1-2 1-3 1-4 4-5 4-8 5-6 5-9 6-7

exact/norm bonds :

1-4 5-6

exact bonds :

1-2 1-3 4-5 4-8 5-9 6-7

Match level :


1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS

L1 STRUCTURE UPLOADED

=> D L1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> S L1

SAMPLE SEARCH INITIATED 14:16:51 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 1948 TO ITERATE

51.3% PROCESSED 1000 ITERATIONS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

0 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

03/14/2005 10807710.trn

BATCH **COMPLETE**
PROJECTED ITERATIONS: 36313 TO 41607
PROJECTED ANSWERS: 0 TO 0

L2 0 SEA SSS SAM L1

=> S L1 SSS FULL
FULL SEARCH INITIATED 14:16:58 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 37690 TO ITERATE

100.0% PROCESSED 37690 ITERATIONS
SEARCH TIME: 00.00.01

4 ANSWERS

L3 4 SEA SSS FUL L1

=> FIL CAPLUS
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
161.33	161.54

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 14:17:04 ON 14 MAR 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 14 Mar 2005 VOL 142 ISS 12
FILE LAST UPDATED: 13 Mar 2005 (20050313/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> S L3
L4

3 L3

=> d 14 ibib abs hitstr tot

L4 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2005 ACS on STN
ACCESSION NUMBER: 2005:141013 CAPLUS
DOCUMENT NUMBER: 142:212335
TITLE: Polyamine and aminothiols compounds and compositions for use in conjunction with cancer therapy
INVENTOR(S): Fahl, William E.; Peebles, Daniel D.; Copp, Richard R.
PATENT ASSIGNEE(S): Wisconsin Alumni Research Foundation, USA
SOURCE: PCT Int. Appl., 142 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005014524	A2	20050217	WO 2004-US26535	20040809
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				


PRIORITY APPLN. INFO.: US 2003-493218P P 20030807

AB The invention provides polyamine and aminothiols compds. and pharmaceutical compns. for administration in conjunction with cancer chemotherapy or radiation therapy. The compds. are administered locally to provide protection against the adverse side-effects of chemotherapy or radiation therapy, such as alopecia, mucositis and dermatitis. Pharmaceutical preps. comprising one or more chemoprotective polyamines or aminothiols formulated for topical or local delivery to epithelial or mucosal cells are disclosed. Methods of administering the pharmaceutical preps. are also disclosed.

IT **844435-75-2**
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (polyamine and aminothiol compds. and compns. for use in conjunction with cancer therapy)

RN 844435-75-2 CAPLUS

CN 2-Butanethiol, 3-(dimethylamino)- (9CI) (CA INDEX NAME)



L4 ANSWER 2 OF 3 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1984:562548 CAPLUS

DOCUMENT NUMBER: 101:162548

TITLE: Substituted cysteamine ligands and their complexes with molybdenum(VI)

AUTHOR(S): Corbin, James L.; Miller, Kenneth F.; Pariyadath, Narayanakutty; Heinecke, Jay; Bruce, Alice E.; Wherland, Scot; Stiefel, Edward I.

CORPORATE SOURCE: Charles F. Kettering Res. Lab., Yellow Springs, OH, 45387, USA

SOURCE: Inorganic Chemistry (1984), 23(21), 3404-12

CODEN: INOCAJ; ISSN: 0020-1669

DOCUMENT TYPE: Journal

LANGUAGE: English

AB New bidentate cysteamine-based ligands containing Me-substituted C and N atoms were prepared Together with known ligands the following complete set has now been prepared: NH₂CH₂CH₂SH, MeNHCH₂CH₂SH, Me₂NCH₂CH₂SH, NH₂C(CH₃)₂CH₂SH,


MeNHC(CH₃)₂CH₂SH, Me₂NC(CH₃)₂CH₂SH, NH₂CH₂C(CH₃)₂SH, MeNHCH₂C(CH₃)₂SH, Me₂NCH₂C(CH₃)₂SH, NH₂C(CH₃)₂C(CH₃)₂SH, MeNHC(CH₃)₂C(CH₃)₂SH, and Me₂NC(CH₃)₂C(CH₃)₂SH. Five of the ligands in this series are new, and their preparation is reported in detail. Also RNHCH₂C(CH₃)₂SH ligands (R = iso-Pr and iso-Bu) are reported for the 1st time. These ligands, LH, were reacted in MeOH with MoO₂(acac)₂ (Hacac = acetylacetone). In most cases MoO₂L₂ resulted. However, in some cases this complex appears to be unstable. The prepns. and spectroscopic properties of the complexes are reported. The low values of ν(Mo-O) for some of the complexes are correlated either with H bonding or with the presence of a skew-trapezoidal-bipyramidal structure. Likewise, electronic absorption spectra differ for complexes with octahedral as opposed to skew-trapezoidal-bipyramidal structures. For a given complex, 170 and 1H NMR spectroscopies are consistent with adoption in solution of the same octahedral or skew-trapezoidal-bipyramidal structure that is found in the solid state. Further, the skew-trapezoidal-bipyramidal complexes display temperature-dependent NMR spectra that are interpreted in terms of configurational averaging probably caused by Mo-N bond cleavage.

IT 91229-44-6P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 91229-44-6 CAPLUS

CN 2-Butanethiol, 3-(dimethylamino)-2,3-dimethyl-, hydrochloride (9CI) (CA INDEX NAME)



● HCl

L4 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1964:447807 CAPLUS

DOCUMENT NUMBER: 61:47807

ORIGINAL REFERENCE NO.: 61:8284a-b

TITLE: Preparation of quaternary ammonium betaine salts

INVENTOR(S): Klass, Donald L.

PATENT ASSIGNEE(S): Pure Oil Co.

SOURCE: 4 pp.

DOCUMENT TYPE: Patent

LANGUAGE: Unavailable

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 3131189		19640428	US	19611016

GI For diagram(s), see printed CA Issue.


AB Carbyl sulfate (I), prepared by the reaction of 2 moles SO₃ and 1 mole ethylene, reacted with a tertiary amine to form betaines. Thus 1.5 g. pyridine (II) in 10 ml. ethylene dichloride was added to 3 g. I in 30 ml. ethylene dichloride (the reaction was exothermic), the liquid decanted from the precipitate, and the precipitate covered with petr. ether and cooled to give

IIa (R = R1 = H), m. 250-5° (HCONMe2). I was also treated with the following to form betaines: quinoline, acridine, trimethylamine, and dimethylaniline (III). Also reported without details were: IIa (R = Ph, R1 = H); Et3NCH2CH2SO3; IIa (R = R1 = Me); and PhNMe2CMe2CMe2SO3. These compds. are useful intermediates for the preparation of detergents. (Cf. U.S. 2,666,788, or Brit. 686,061.)

IT 97176-62-0, Ammonium, dimethylphenyl(1,1,2-trimethyl-2-sulfopropyl), hydroxide, inner salt
(preparation of)

RN 97176-62-0 CAPLUS

CN Dimethylphenyl(1,1,2-trimethyl-2-sulfopropyl)ammonium hydroxide, inner salt (7CI) (CA INDEX NAME)



=> FIL REGISTRY

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

18.42

179.96

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-2.19

-2.19

FILE 'REGISTRY' ENTERED AT 14:21:56 ON 14 MAR 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 13 MAR 2005 HIGHEST RN 845467-46-1

DICTIONARY FILE UPDATES: 13 MAR 2005 HIGHEST RN 845467-46-1

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.



Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:

<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10807710a.str



chain nodes :
 1 2 3 4 5 6 7 8 9
 chain bonds :
 1-2 1-3 1-4 4-5 4-8 5-6 5-9 6-7
 exact/norm bonds :
 1-4 4-8 5-6 5-9 6-7
 exact bonds :
 1-2 1-3 4-5

G1: Ak, Ph

G2: H, CH3

Match level :


1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS

L5 STRUCTURE UPLOADED

=> d 15

L5 HAS NO ANSWERS

L5 STR



G1 Ak, Ph

G2 H, Me

Structure attributes must be viewed using STN Express query preparation.

=> s 15

SAMPLE SEARCH INITIATED 14:22:20 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 5592 TO ITERATE

03/14/2005 10807710.trn

17.9% PROCESSED 1000 ITERATIONS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

0 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 107357 TO 116323
PROJECTED ANSWERS: 0 TO 0

L6 0 SEA SSS SAM L5

=> s 15 sss full
FULL SEARCH INITIATED 14:22:26 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 109275 TO ITERATE

100.0% PROCESSED 109275 ITERATIONS
SEARCH TIME: 00.00.02

36 ANSWERS

L7 36 SEA SSS FUL L5

=> FIL CAPLUS
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
161.33	341.29

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
0.00	-2.19

CA SUBSCRIBER PRICE

FILE 'CAPLUS' ENTERED AT 14:22:37 ON 14 MAR 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 14 Mar 2005 VOL 142 ISS 12
FILE LAST UPDATED: 13 Mar 2005 (20050313/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 17
L8

18 L7

=> FIL REGISTRY
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION

03/14/2005 10807710.trn

FULL ESTIMATED COST	1.80	343.09
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-2.19

FILE 'REGISTRY' ENTERED AT 14:24:46 ON 14 MAR 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 13 MAR 2005 HIGHEST RN 845467-46-1
DICTIONARY FILE UPDATES: 13 MAR 2005 HIGHEST RN 845467-46-1

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005


Please note that search-term pricing does apply when
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more
information enter HELP PROP at an arrow prompt in the file or refer
to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10807710b.str



chain nodes :

2 3 4 5 6 7

ring nodes :

1 11 12 13 14 15

chain bonds :

1-2 2-3 2-6 3-4 3-7 4-5

ring bonds :

1-11 1-12 11-14 12-13 13-15 14-15

exact/norm bonds :

1-2 1-11 1-12 2-6 3-4 3-7 4-5 11-14 12-13 13-15 14-15

exact bonds :

2-3

03/14/2005 10807710.trn

isolated ring systems :
containing 1 :

G1: Ak, Ph

G2: H, CH3

Match level :


1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 11:CLASS 12:CLASS
13:CLASS 14:CLASS 15:CLASS

L9 STRUCTURE UPLOADED

=> d 19

L9 HAS NO ANSWERS

L9 STR



G1 Ak, Ph

G2 H, Me

Structure attributes must be viewed using STN Express query preparation.

=> s 19

SAMPLE SEARCH INITIATED 14:25:15 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 3163 TO ITERATE

31.6% PROCESSED 1000 ITERATIONS

INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)

SEARCH TIME: 00.00.01

0 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 59887 TO 66633

PROJECTED ANSWERS: 0 TO 0

03/14/2005 10807710.trn

L10 0 SEA SSS SAM L9

=> s l9 sss full
FULL SEARCH INITIATED 14:25:23 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 62510 TO ITERATE

100.0% PROCESSED 62510 ITERATIONS
SEARCH TIME: 00.00.01

17 ANSWERS

L11 17 SEA SSS FUL L9

=> FIL CAPLUS
~~COST IN U.S. DOLLARS~~

FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
161.33	504.42

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

CA SUBSCRIBER PRICE

SINCE FILE	TOTAL
ENTRY	SESSION
0.00	-2.19

FILE 'CAPLUS' ENTERED AT 14:25:32 ON 14 MAR 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 14 Mar 2005 VOL 142 ISS 12
FILE LAST UPDATED: 13 Mar 2005 (20050313/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d his

(FILE 'HOME' ENTERED AT 14:16:29 ON 14 MAR 2005)

FILE 'REGISTRY' ENTERED AT 14:16:38 ON 14 MAR 2005

L1 STRUCTURE UPLOADED
L2 0 S L1
L3 4 S L1 SSS FULL

FILE 'CAPLUS' ENTERED AT 14:17:04 ON 14 MAR 2005

L4 3 S L3

FILE 'REGISTRY' ENTERED AT 14:21:56 ON 14 MAR 2005

L5 STRUCTURE UPLOADED
L6 0 S L5
L7 36 S L5 SSS FULL

10807710.trn

Page 12

14:27

FILE 'CAPLUS' ENTERED AT 14:22:37 ON 14 MAR 2005
L8 18 S L7

FILE 'REGISTRY' ENTERED AT 14:24:46 ON 14 MAR 2005
L9 STRUCTURE UPLOADED
L10 0 S L9
L11 17 S L9 SSS FULL

FILE 'CAPLUS' ENTERED AT 14:25:32 ON 14 MAR 2005

=> s l11

L12 20 L11

=> s l8 and py<=2002

22589460 PY<=2002

L13 17 L8 AND PY<=2002

=> s l12 and py<=2002

22589460 PY<=2002

L14 17 L12 AND PY<=2002

=> d l13 ibib abs hitstr tot

L13 ANSWER 1 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1999:671830 CAPLUS

DOCUMENT NUMBER: 132:35935

TITLE: Diastereomeric sulfinates derived from
(L)-N-methylephedrine: synthesis, applications and
rearrangements

AUTHOR(S): Drabowicz, Jozef; Bujnicki, Bogdan; Biscarini, Paolo;
Mikolajczyk, Marian

CORPORATE SOURCE: Centre of Molecular and Macromolecular Studies, Polish
Academy of Sciences, Lodz, 90-363, Pol.

SOURCE: Tetrahedron: Asymmetry (1999), 10(16),
3177-3187

CODEN: TASYE3; ISSN: 0957-4166

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 132:35935

AB The reaction of sulfinyl chlorides with (L)-N-methylephedrine alone or in
the presence of tertiary amines was found to produce diastereomeric
sulfinates with diastereomeric purities up to 90%. The diastereomeric
ratio is strongly influenced by the nature of substituents on the sulfinyl
chlorides and to some extent by the reaction conditions. In a few cases,
the pure diastereomers were isolated by chromatog. and used for the preparation
of optically active sulfoxides. The silica gel catalyzed rearrangement of
sulfinates to the corresponding sulfones is also discussed.


IT 252230-36-7P

RL: SPN (Synthetic preparation); PREP (Preparation)
(diastereomeric sulfinates derived from (L)-N-methylephedrine,
synthesis, applications and rearrangements)

RN 252230-36-7 CAPLUS

CN Benzeneethanamine, N,N, α -trimethyl- β -(methylsulfonyl)-,
(α S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 2 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1995:809918 CAPLUS
 DOCUMENT NUMBER: 123:286349
 TITLE: Thiol and disulfide derivatives of Ephedra alkaloids.
 2. A mechanistic study of their effect on the addition
 of diethylzinc to benzaldehyde
 AUTHOR(S): Fitzpatrick, Kevin; Hulst, Ron; Kellogg, Richard M.
 CORPORATE SOURCE: Dep. of Organic and Molecular Inorganic Chem., Univ.
 of Groningen, Groningen, 9747 AG, Neth.
 SOURCE: Tetrahedron: Asymmetry (1995), 6(8), 1861-4
 CODEN: TASYE3; ISSN: 0957-4166
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 123:286349
 AB Non-linear correlations between the ee of the product, PhCH(OH)Et, and the
 catalyst, ephedrine thiol or disulfide derivs., were observed Osmotic
 measurements indicate a high degree of aggregation of the Zn thiolates.
 The behavior of the thiol derivs. deviates sharply from that of
 N-methylephedrine.
 IT 2218-21-5
 RL: CAT (Catalyst use); USES (Uses)
 (effect of thiol and disulfide derivs. of ephedrine on addition of Et2Zn
 to benzaldehyde)
 RN 2218-21-5 CAPLUS
 CN Benzenemethanethiol, α -[1-(dimethylamino)ethyl]-, hydrochloride,
 [R-(R*,S*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl

L13 ANSWER 3 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1990:531411 CAPLUS
 DOCUMENT NUMBER: 113:131411
 TITLE: Concerning mass spectra of β -heterosubstituted
 thioethers

AUTHOR(S): Carreno, M. Carmen; Carretero, Juan C.; Garcia Ruano, Jose L.; Martinez, M. Carmen
 CORPORATE SOURCE: Dep. Quim., Univ. Auton. Madrid, Madrid, 28049, Spain
 SOURCE: Organic Mass Spectrometry (1990), 25(6), 339-42

CODEN: ORMSBG; ISSN: 0030-493X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The mass spectra of 32 erythro- and threo-RCHXCHRSMe (R = Me, Ph; X = F, Cl, Br, iodo, OH, OMe, OAc, NH₂, NMe₂) were interpreted in detail. Linear free-energy relationships were determined between the ratio of the relative abundances of the RCHX⁺ and RCHSMe⁺ peaks and the inductive and resonance σ consts. of X.

IT 97241-41-3 97241-45-7 97241-46-8

97241-47-9


RL: PRP (Properties)

(mass spectrum of)

RN 97241-41-3 CAPLUS

CN 2-Butanamine, N,N-dimethyl-3-(methylthio)-, (R*,S*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 97241-45-7 CAPLUS

CN 2-Butanamine, N,N-dimethyl-3-(methylthio)-, (R*,R*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 97241-46-8 CAPLUS

CN Benzeneethanamine, N,N-dimethyl- β -(methylthio)- α -phenyl-, (R*,S*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 97241-47-9 CAPLUS

CN Benzeneethanamine, N,N-dimethyl- β -(methylthio)- α -phenyl-,
(R*,R*)- (9CI) (CA INDEX NAME)

Relative stereochemistry.



L13 ANSWER 4 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1990:458617 CAPLUS

DOCUMENT NUMBER: 113:58617

TITLE: Reactions of organic anions. 168. Reactions of
2-(dialkylamino)arylacetonitriles with acetylenes
under basic conditions. A simple synthesis of
substituted mono- and diketones

AUTHOR(S): Zdrojewski, T.; Jonczyk, A.

CORPORATE SOURCE: Dep. Chem., Tech. Univ., Warsaw, PL-00-662, Pol.

SOURCE: Synthesis (1990), (3), 224-33
CODEN: SYNTBF; ISSN: 0039-7881

DOCUMENT TYPE: Journal

LANGUAGE: English


OTHER SOURCE(S): CASREACT 113:58617

AB The reaction of $\text{RCH}(\text{CN})\text{NR}_{12}$ (I; R = Ph, 4-MeC₆H₄, 4-MeOC₆H₄; R₁ = Me; NR₁₂ = piperidino, morpholino, etc.) with $\text{R}_2\text{C}.\text{tp}.\text{bond}.\text{CH}$ (II; R₂ = Ph, MeS) gave $\text{R}_{12}\text{NCR}(\text{CN})\text{CH}:\text{CHR}_2$ (III) and/or $\text{R}_{12}\text{NCR}(\text{CN})\text{CHR}_2\text{CH}:\text{CRNR}_{12}$; the product depended on the basicity of the amino group in III. I also added to C-1 of II (R₂ = EtO) to give $\text{R}_{12}\text{NCR}(\text{CN})\text{C}(\text{OEt})\text{:CH}_2$. All these products could be hydrolyzed to give mono- or diketones.

IT 128407-41-0P 128407-44-3P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(preparation and hydrolysis of)

RN 128407-41-0 CAPLUS

CN Benzeneacetonitrile, α -(dimethylamino)- α -[3-(dimethylamino)-3-(4-methylphenyl)-1-(methylthio)-2-propenyl]-4-methyl- (9CI) (CA INDEX NAME)

RN 128407-44-3 CAPLUS

CN Benzeneacetonitrile, α -(dimethylamino)-4-methyl- α -[1-(methylthio)-3-phenyl-3-(1-piperidiny)-2-propenyl]- (9CI) (CA INDEX NAME)




IT 128407-50-1P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 128407-50-1 CAPLUS

CN Benzenepentanenitrile, α-(dimethylamino)-4-methyl-α-(4-methylphenyl)-β-(methylthio)-δ-oxo- (9CI) (CA INDEX NAME)



L13 ANSWER 5 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1985:505278 CAPLUS

DOCUMENT NUMBER: 103:105278

TITLE: L-Cysteine-, L-methionine- and D-penicillamine-derived ligands for transition metal catalyzed carbon-carbon bond-forming reactions

AUTHOR(S): Griffin, John H.; Kellogg, Richard M.

CORPORATE SOURCE: Dep. Org. Chem., Univ. Groningen, Groningen, 9747 AG, Neth.

SOURCE: Journal of Organic Chemistry (1985), 50(18), 3261-6

CODEN: JOCEAH; ISSN: 0022-3263

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 103:105278


AB L-Cysteine, L-methionine, and D-penicillamine were converted to their corresponding S-alkylated dimethylamino alcs. by alkylation of sulfur and nitrogen followed by reduction of the carboxyl group. One-pot mesylation followed by treatment with lithium diphenylphosphide provides the corresponding phosphine derivative, 2-(dimethylamino)-1-(diphenylphosphino)-3-(methylthio)propane (cysphos), the S-iso-Pr derivative (isopropylcycphos), 2-(dimethylamino)-1-(diphenylphosphino)-4-(methylthio)butane (methphos), and 2-(dimethylamino)-1-(diphenylphosphino)-3-methyl-3-(methylthio)butane (penphos). The latter ligand had an enantiomeric excess of 82%, whereas the other ligands are enantiomerically pure. These compds. were used as ligands for the solubilization of catalytic amts. of NiCl₂, which is used to mediate the coupling of vinyl bromide with the (racemic) Grignard reagent or 1-phenyl-1-chloroethane to give 3-phenyl-1-butene in excellent yields and with enantiomeric excesses of up to 65% (methphos). The cross coupling of vinyl bromide to the Grignard reagent of 2-octyl chloride was also examined; enantiomeric excesses of 3-methyl-1-nonene of up to 18% are obtained (penphos).

IT 97071-94-8P 97072-02-1P

RL: SPN (Synthetic preparation); PREP (Preparation)


(preparation of)
 RN 97071-94-8 CAPLUS
 CN 1-Butanol, 2-(dimethylamino)-3-methyl-3-(methylthio)-, (R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 97072-02-1 CAPLUS
 CN Benzeneacetic acid, α -methoxy- α -(trifluoromethyl)-, 2-(dimethylamino)-3-methyl-3-(methylthio)butyl ester, [S-(R*,S*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 97071-99-3P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, as ligand for asym. nickel-catalyzed crosscoupling of Grignard reagents with vinyl bromide)
 RN- 97071-99-3 CAPLUS
 CN 2-Butanamine, 1-(diphenylphosphino)-N,N,3-trimethyl-3-(methylthio)-, (R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 6 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1985:437072 CAPLUS
 DOCUMENT NUMBER: 103:37072
 TITLE: Syntheses of β -heterosubstituted thioethers
 AUTHOR(S): Carretero, J. Carlos; Garcia-Ruano, Jose L.; Martinez, M. Carmen; Rodriguez, Jesus H.
 CORPORATE SOURCE: Dep. Quim. Org., Univ. Auton. Madrid, Madrid, 28049, Spain
 SOURCE: Journal of Chemical Research, Synopses (1985), (1), 6-7

CODEN: JRPSDC; ISSN: 0308-2342

DOCUMENT TYPE:

Journal

LANGUAGE:

English

OTHER SOURCE(S):

CASREACT 103:37072

AB The thio ethers erythro- and threo-RCHR₁CHRSMe (I; R = Me, Ph; R₁ = OH, OMe, F, Cl, Br, iodo, NH₂, NMe₂) were prepared starting from readily available substrates. I (R = Me, R₁ = OH) were readily prepared by treatment of the corresponding MeCHBrCHMeOH with NaSMe in MeOH at 0° for 3 h; the reaction occurs with retention of configuration. Treatment of I (R = Me, R₁ = OH) with HCl(g) in CHCl₃ at room temperature gave the corresponding I (R = Me, R₁ = Cl).

IT 97241-41-3P 97241-45-7P 97241-46-8P


97241-47-9P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 97241-41-3 CAPLUS

CN 2-Butanamine, N,N-dimethyl-3-(methylthio)-, (R*,S*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 97241-45-7 CAPLUS

CN 2-Butanamine, N,N-dimethyl-3-(methylthio)-, (R*,R*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 97241-46-8 CAPLUS

CN Benzeneethanamine, N,N-dimethyl-β-(methylthio)-α-phenyl-,
(R*,S*)- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 97241-47-9 CAPLUS

CN Benzeneethanamine, N,N-dimethyl-β-(methylthio)-α-phenyl-,

(R*,R*)- (9CI) (CA INDEX NAME)

Relative stereochemistry.



L13 ANSWER 7 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1984:562548 CAPLUS

DOCUMENT NUMBER: 101:162548

TITLE: Substituted cysteamine ligands and their complexes with molybdenum(VI)

AUTHOR(S): Corbin, James L.; Miller, Kenneth F.; Pariyadath, Narayanakutty; Heinecke, Jay; Bruce, Alice E.; Wherland, Scot; Stiefel, Edward I.

CORPORATE SOURCE: Charles F. Kettering Res. Lab., Yellow Springs, OH, 45387, USA

SOURCE: Inorganic Chemistry (1984), 23(21), 3404-12

CODEN: INOCAJ; ISSN: 0020-1669

DOCUMENT TYPE: Journal

LANGUAGE: English


AB New bidentate cysteamine-based ligands containing Me-substituted C and N atoms were prepared. Together with known ligands the following complete set has now been prepared: $\text{NH}_2\text{CH}_2\text{CH}_2\text{SH}$, $\text{MeNHCH}_2\text{CH}_2\text{SH}$, $\text{Me}_2\text{NCH}_2\text{CH}_2\text{SH}$, $\text{NH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{SH}$, $\text{MeNHC}(\text{CH}_3)_2\text{CH}_2\text{SH}$, $\text{Me}_2\text{NC}(\text{CH}_3)_2\text{CH}_2\text{SH}$, $\text{NH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{SH}$, $\text{MeNHCH}_2\text{C}(\text{CH}_3)_2\text{SH}$, $\text{Me}_2\text{NCH}_2\text{C}(\text{CH}_3)_2\text{SH}$, $\text{NH}_2\text{C}(\text{CH}_3)_2\text{C}(\text{CH}_3)_2\text{SH}$, $\text{MeNHC}(\text{CH}_3)_2\text{C}(\text{CH}_3)_2\text{SH}$, and $\text{Me}_2\text{NC}(\text{CH}_3)_2\text{C}(\text{CH}_3)_2\text{SH}$. Five of the ligands in this series are new, and their preparation is reported in detail. Also $\text{RNHCH}_2\text{C}(\text{CH}_3)_2\text{SH}$ ligands ($\text{R} = \text{iso-Pr}$ and iso-Bu) are reported for the 1st time. These ligands, LH , were reacted in MeOH with $\text{MoO}_2(\text{acac})_2$ ($\text{Hacac} = \text{acetylacetone}$). In most cases MoO_2L_2 resulted. However, in some cases this complex appears to be unstable. The preps. and spectroscopic properties of the complexes are reported. The low values of $\nu(\text{Mo-O})$ for some of the complexes are correlated either with H bonding or with the presence of a skew-trapezoidal-bipyramidal structure. Likewise, electronic absorption spectra differ for complexes with octahedral as opposed to skew-trapezoidal-bipyramidal structures. For a given complex, ^{17}O and ^1H NMR spectroscopies are consistent with adoption in solution of the same octahedral or skew-trapezoidal-bipyramidal structure that is found in the solid state. Further, the skew-trapezoidal-bipyramidal complexes display temperature-dependent NMR spectra that are interpreted in terms of configurational averaging probably caused by Mo-N bond cleavage.

IT 91229-44-6P

RL: SPN (Synthetic preparation); PREP (Preparation of)

RN 91229-44-6 CAPLUS

CN 2-Butanethiol, 3-(dimethylamino)-2,3-dimethyl-, hydrochloride (9CI) (CA INDEX NAME)



● HCl

L13 ANSWER 8 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1982:216922 CAPLUS

DOCUMENT NUMBER: 96:216922

TITLE: Thioammonium ions. Azasulfenylation reactions

AUTHOR(S): Caserio, Marjorie C.; Kim, Jhong K.

CORPORATE SOURCE: Dep. Chem., Univ. California, Irvine, CA, 92717, USA

SOURCE: Journal of the American Chemical Society (1982), 104(11), 3231-3

CODEN: JACSAT; ISSN: 0002-7863

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 96:216922

AB Attempts to prepare elusive (alkylthio)ammonium ions have given evidence for their formation as reactive intermediates in alkylation of sulfenamides and in the sulfenylation of amines. In the presence of alkenes, (alkylthio)ammonium ions are trapped as 1-ammonium-2-thio adducts formed by anti-addition. These same adducts were obtained more conveniently in 2 steps: addition of (alkylthio)sulfonium ions to the alkene, followed by displacement of the sulfonium group by an amine nucleophile. The displacement step proceeds by the reversible dissociation of 1-sulfonium-2-thio adducts to episulfonium ions. The displacement reaction is general because the episulfonium ion can be trapped by a variety of nucleophiles.

IT 81206-95-3P 81206-97-5P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 81206-95-3 CAPLUS


CN 2-Butanaminium, N,N,N-trimethyl-3-(methylthio)-, (R*,R*)-, tetrafluoroborate(1-) (9CI) (CA INDEX NAME)

CM 1

CRN 81206-94-2


CMF C8 H20 N S

Relative stereochemistry.



CM 2

CRN 14874-70-5
CMF B F4
CCI CCS




RN 81206-97-5 CAPLUS
CN 2-Butanaminium, N,N,N-trimethyl-3-(methylthio)-, (R*,S*)-, tetrafluoroborate(1-) (9CI) (CA INDEX NAME)

CM 1


CRN 81206-96-4
CMF C8 H20 N S

Relative stereochemistry.



CM 2


CRN 14874-70-5
CMF B F4
CCI CCS



L13 ANSWER 9 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
ACCESSION NUMBER: 1979:574952 CAPLUS
DOCUMENT NUMBER: 91:174952
TITLE: Thermolysis of Mannich bases from β -oxo sulfoxides, benzaldehyde and secondary amines
AUTHOR(S): Boehme, Horst; Clement, Bernd
CORPORATE SOURCE: Pharm.-Chem. Inst., Philipps-Univ., Marburg, 355, Fed. Rep. Ger.
SOURCE: Archiv der Pharmazie (Weinheim, Germany) (1979), 312(6), 531-4

DOCUMENT TYPE:
LANGUAGE:
GI

CODEN: ARPMAS; ISSN: 0365-6233
Journal
German



AB Stable Mannich bases I [R = Me, R₂ = (CH₂)₄, CH₂CH₂OCH₂CH₂] were obtained as mixts. of 4 diastereoisomeric forms, stipulated by the 3 chiral centers, by condensation of PhCOCH₂S(O)Me with PhCHO and HNR₂. Amine elimination occurred on heating I (R = Me) >180° to give a single (E) diastereomer of the propenone II.


IT 71679-36-2P 71698-77-6P 71698-78-7P
71698-79-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and thermolysis of)

RN 71679-36-2 CAPLUS

CN 1-Propanone, 3-(dimethylamino)-2-(methylsulfinyl)-1,3-diphenyl-,
[2R*(R*),3R*]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 71698-77-6 CAPLUS

CN 1-Propanone, 3-(dimethylamino)-2-(methylsulfinyl)-1,3-diphenyl-,
[2R*(S*),3R*]- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 71698-78-7 CAPLUS

CN 1-Propanone, 3-(dimethylamino)-2-(methylsulfinyl)-1,3-diphenyl-,

[2R*(R*),3S*]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 71698-79-8 CAPLUS

CN 1-Propanone, 3-(dimethylamino)-2-(methylsulfinyl)-1,3-diphenyl-,
[2R*(S*),3S*]- (9CI) (CA INDEX NAME)

Relative stereochemistry.



L13 ANSWER 10 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1979:97354 CAPLUS

DOCUMENT NUMBER: 90:97354

TITLE: Studies on chemical protectors against radiation. XX.
Radioprotective mechanism of phenethylamine compounds

AUTHOR(S): Shinoda, Masato; Onoda, Makoto; Sato, Kazunori;
Nemoto, Kyuichi; Kamogawa, Asahi

CORPORATE SOURCE: Hoshi Coll. Pharm., Tokyo, Japan


SOURCE: Yakugaku Zasshi (1978), 98(11), 1512-17

CODEN: YKKZAJ; ISSN: 0031-6903

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

GI



I, R¹ and R² = H, OH, SH, SSO₃H, or SC(:NH)NH₂,

R³ and R⁴ = H or Me, R⁵ = H, Ac, alkyl, NH₂,

or Bz, R⁶ and R⁷ = H or OH

AB Correlation between radioprotection by phenethylamines (I) and anoxia was

examined from their action on femoral artery blood flow and on blood pressure of the rat common carotid and isolated dorsal arteries. The radioprotection mechanism by vasoconstriction-anoxia was present in the phenethylamine-type and ephedrine-type compds., and possible participation of anoxia was suggested in some of S-ephedrine- and isothioureia-type compds. In disulfide- and thiosulfate-type compds., mechanisms other than anoxia were considered to be present.


IT 942-48-3

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
(radioprotective activity of)

RN 942-48-3 CAPLUS

CN Benzenemethanethiol, α -[1-(dimethylamino)ethyl]-, [S-(R*,R*)]- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



L13 ANSWER 11 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1978:31877 CAPLUS

DOCUMENT NUMBER: 88:31877

TITLE: Studies on chemical protectors against radiation.
XVII. Radioprotective activities of phenethylamine compounds

AUTHOR(S): Shinoda, Masato; Ohta, Setsuko; Takagi, Yoshinari

CORPORATE SOURCE: Hoshi Coll. Pharm., Tokyo, Japan

SOURCE: Yakugaku Zasshi (1977), 97(10), 1117-24

CODEN: YKKZAJ; ISSN: 0031-6903

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

AB The relation between chemical structure and radioprotective activity was examined with 80 phenethylamine compds. and various amines. A strong radioprotective effect was shown by phenethylamine-HCl [156-28-5] and by tyramine-HCl [60-19-5], dopamine-HCl [62-31-7], norepinephrine [51-41-2], and epinephrine-HCl [55-31-2], which have a phenolic hydroxyl in their mol., but the corresponding amino acids were ineffective. Only a weak effect was shown by the ephedrine isomers, but a markedly strong effect was shown by compds. with a side chain substituted with SH, isothioureia, or thiosulfuric acid, and by the S-S ephedrine compds. Comparison of the isomers of these compds. showed that the L-erythro type compds. were more effective.


IT 942-50-7

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
(radioprotective activity of)

RN 942-50-7 CAPLUS

CN Benzenemethanethiol, α -[1-(dimethylamino)ethyl]-, hydrochloride,
[S-(R*,R*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl

L13 ANSWER 12 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1965:410290 CAPLUS

DOCUMENT NUMBER: 63:10290

ORIGINAL REFERENCE NO.: 63:1821h,1822f-h,1823a-c

TITLE: Potential radiation-protective agents. III. Mercapto analogs related to ephedrine

AUTHOR(S): Bhat, K. Venkatramana; McCarthy, Walter C.

CORPORATE SOURCE: Univ. of Washington, Seattle

SOURCE: Journal of Pharmaceutical Sciences (1965), 54(2), 225-8

CODEN: JPMSAE; ISSN: 0022-3549

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 63:10290

AB cf. CA 62, 5214f. Since certain sympathomimetic amines have been shown to possess radiation protective activity, the mercapto analogs related to ephedrine were selected for investigation. (+)-PhCHClCHMeNHMe.HCl (I.HCl) (19 g.) and 21.4 g. Na₂S₂O₃·5H₂O dissolved in 40 ml. hot H₂O and the solution let stand 1 hr., heated 2 hrs. on a water bath, and cooled gave 20.5 g. Bunte salt PhCH(S₂O₃⁻)CHMeN+H₂Me (II), m. 186-7° (H₂O), [α]₂₅^D 221° (c 1, H₂O). To 22 g. (+)-I.HCl in 150 ml. anhydrous MeOH was added 23 ml. MeOH-MeONa (10% Na weight/volume), followed by MeCOSNa solution [prepared from 7.5 ml. MeCOSH and 23 ml. MeOH-MeONa (10% Na weight/volume)], the mixture refluxed 3 hrs. and filtered, and the filtrate evaporated in vacuo (rotary evaporator) to give 9.5 g. (+)-PhCH(SH)CHMeNMeAc (III), m. 112-13° (dilute EtOH), [α]₂₅^D 190° (c 1, EtOH). Oxidation of (+)-III with iodine gave (+)-(AcMeNCHMeCHPhS)₂ (IV), m. 172-3° (dilute EtOH), [α]₂₅^D 390° (c 1, 95% EtOH). (+)-III (5.6 g.) in 50 ml. EtOH refluxed 3 hrs. with 25 ml. concentrated HCl

and

the solution evaporated in vacuo (rotary evaporator) gave 4.5 g. unchanged (+)-III, m. 112-13° (dilute EtOH). (+)-III (5.6 g.) in 100 ml. 20% aqueous NaOH refluxed 24 hrs., cooled, and saturated with CO₂ gave [by air

oxidation

of (+)-III] 3.2 g. (+)-IV, m. 172-3° (dilute EtOH). (+)-IV (4.5 g.)

extracted from the thimble of a Soxhlet extractor by a refluxing solution of

1.8

g. LiAlH₄ in 275 ml. Et₂O [after 72 hrs. all (+)-IV had been extracted], 0.9 ml. H₂O added dropwise with stirring, then 4.9 ml. concentrated HCl, the

precipitate

filtered off, extracted with Et₂O, the extract filtered, the combined filtrates dried and evaporated in vacuo (rotary evaporator), and the semisolid residue recrystd. (EtOH) gave 2 g. (EtMeNCHMeCHPhS)₂, m. 160-1°, [α]₂₅^D 715° (c 1, C₆H₆). (+)-I.HCl (22 g.) in 200 ml. absolute


EtOH treated with 46 ml. EtOH-EtONa (5% Na weight/volume), followed by 16 g. EtOCS₂K, and the mixture heated 3 hrs. on a steam bath and worked up gave 6

g. (-)-3,4-dimethyl-5-phenylthiazolidine-2-thione (V), m. 65-6° (iso-Pr₂O), [α]_{25D} -155° (c 1, EtOH), which was resistant to hydrolysis by boiling 48% HBr. (-)-V (8 g.) in 200 ml. anhydrous Et₂O added dropwise to 3.79 g. LiAlH₄ in 100 ml. Et₂O and the solution refluxed 90 hrs., hydrolyzed by adding dropwise 7.2 ml. H₂O and 9.7 ml. concentrated HCl, and worked up gave 4.5 g. (+)-PhCH(SH)CHMeNMe₂, m. 75-6° (absolute EtOH), [α]_{25D} 150° (c 1, 0.1N HCl), which oxidized in EtOH with 1% iodine solution gave (+)-(Me₂NCHMeCHPhS)₂, m. 196-7° (EtOH), [α]_{25D} 300° (c 1, 0.1N HCl). Chloral (15 g.) added dropwise to 16.5 g. (-)-ephedrine (VI) with stirring and ice cooling and the mixture stirred 2 hrs. at room temperature and concentrated in vacuo (rotary evaporator) gave a mixture of oil and crystalline solid; recrystn. of the solid from absolute EtOH gave 1 g. (-)-VI.HCl, m. 217-18°; distillation of the oil gave 6 g. N-formyl-(-)-ephedrine (VII), b₁ 168-74°, ν 1665 cm.⁻¹ Excess SOCl₂ added dropwise to VII with stirring and ice cooling and the mixture stirred 1.5 hrs. at room temperature and worked up gave (+)-pseudoephedrine-HCl (VIII.HCl), m. 182-3°, [α]_{25D} 60° (c 1, H₂O). SOCl₂ (1.13 ml.) added dropwise to an ice cold solution of 2.07 g. N-acetylephephedrine in 100 ml. CHCl₃ and the mixture stirred 30 min. at room temperature, heated 30 min. on a water bath, and worked up gave 1.5 g. O-acetyl-(+)-pseudo-ephedrine-HCl, m. 185-6° (iso-PrOH), [α]_{25D} 98° (c 1, H₂O). A mixture of 10.7 g. (-)-VI.H₂SO₄ and 12 g. 4-MeC₆H₄SO₂Cl (IX) shaken with 60 ml. 10% aqueous NaOH and cooled in ice gave 13 g. N-p-tolylsulfonyl-(-)-ephedrine (X), m. 117-18° (60% EtOH), [α]_{25D} -27.5° (c 1, 95% EtOH), ν 3570 cm.⁻¹ Similar tosylation of VIII.HCl gave 84% N-p-tolylsulfonyl-(+)-pseudo-ephedrine, m. 93-4.5°, [α]_{25D} 25° (c 1, 95% EtOH). X (3.2 g.) and 2.5 g. IX in 8 ml. C₅H₅N stirred 4 hrs. at room temperature and worked up gave 94% unchanged X, m. 117-18°. When the mixture was heated at 80°, unchanged X was also recovered. Refluxing the mixture gave an oil, which could not be crystallized Neither II, when administered to mice in a dosage of 51-150 mg./kg., nor 3,4-dimethyl-5-phenyl-2-iminothiazolidine-HCl (McCarthy and Ho, CA 56, 8612f), when administered to mice in a dose of 50 mg./kg, or less, showed protective activity against radiation.

IT 91341-40-1, α-Toluenethiol, α-[1-(dimethylamino)ethyl]-, (+)- (preparation of)

RN 91341-40-1 CAPLUS

CN α-Toluenethiol, α-[1-(dimethylamino)ethyl]- (6CI, 7CI) (CA INDEX NAME)



L13 ANSWER 13 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1965:74011 CAPLUS
 DOCUMENT NUMBER: 62:74011
 ORIGINAL REFERENCE NO.: 62:13088a-d
 TITLE: 1-Substituted 2-aminoethane derivatives
 INVENTOR(S): Takamatsu, Hideji; Nishimura, Haruki
 PATENT ASSIGNEE(S): Dainippon Pharmaceutical Co., Ltd.
 SOURCE: 2 pp.
 DOCUMENT TYPE: Patent

LANGUAGE: Unavailable
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 39022953		19641015	JP	19610415 <--


AB A mixture of 4 g. erythro-1-phenyl-1-dimethylamino-2-chloropropane-HCl (I) and 5.9 g. K thioacetate is refluxed in 60 cc. MeOH 30 min. to give 2.0 g. erythro-1-phenyl-1-mercapto-2-dimethylaminopropane (Ia), b7 109-12°. I (1.9 g.) is refluxed in 30 cc. 10% NaOH 2.5 hrs. to give 1.1 g. erythro-1-phenyl-2-dimethylaminopropanol (hydrochloride m. 206-7°). I (3.3 g.) is heated with 3.55 g. Na2S2O3, in 10 cc. H2O with steam 15 min. to give erythro-1-phenyl-2-dimethylaminopropane-1-thiosulfuric acid (II) m. 207-8° (decomposition). II (3 g.) is refluxed in 30 cc. 10% HCl 1.5 hrs., the product alkalized with Na2CO3 and extracted with ether to give 1.8 g. Ia, b3 80°. Similarly, other compds. prepared were (starting material, reagent and yield, name, and m.p. of the product given): 1-phenyl-1-dimethylamino-2-chloroethane-HCl (III) 1.5 g., K thioacetate, 1.2 g. 1-phenyl-1-mercapto-2-dimethylaminoethane-HCl, 190° (decomposition); III 8 g., Na2S2O3, 8 g., 1-phenyl-2-dimethylaminoethane-1-thiosulfuric acid, 208° (decomposition); III 2 g., Na2S reagent (7.2 g. Na2S + 10 cc. H2O + 20 cc. MeOH saturated with H2S) (reaction condition heated with steam 30 min.), 1.1 g., 1-phenyl-1-mercapto-2-dimethylaminoethane hydrochloride, 191° (decomposition); III 1 g., 10% NaOH, 0.3 g., 1-phenyl-2-dimethylaminoethanol-HCl, --; threo-1-phenyl-1-methylamino-2-chloropropane-HCl (IV) 1 g., Na2S reagent, 0.6 g., threo-1-phenyl-1-mercapto-2-methylaminopropane, -- (b2 93-6°); IV 5 g., Na2S2O3, 4 g., threo-1-phenyl-2-methylaminopropane-1-thiosulfuric acid (V), 190° (decomposition); erythro isomer of IV 1.85 g., Na2S2O3, 1.7 g. erythro isomer of V, 239° (decomposition); erythro isomer of IV 1.3 g., K thioacetate, acetyl derivative (VI) of erythro-1-phenyl-1-mercapto-2-methylaminopropane (VII), -- (VI is heated with 20% HCl in a sealed tube at 140-5° 3 hrs. to give 0.4 g. VII, b6 104°). These compds. are useful as antispasmodics.

IT 942-47-2, α -Toluenethiol, α -[1-(dimethylamino)ethyl]-, erythro- (preparation of)

RN 942-47-2 CAPLUS

CN α -Toluenethiol, α -[1-(dimethylamino)ethyl]-, erythro- (8CI)
 (CA INDEX NAME)


Relative stereochemistry.



L13 ANSWER 14 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1965:8905 CAPLUS
 DOCUMENT NUMBER: 62:8905
 ORIGINAL REFERENCE NO.: 62:1588c-e
 TITLE: Phenylmercaptoalkylamines. IV. Configurations of
 N-alkyl-1-phenyl-2-chloropropylamines and their

rearrangements with nucleophilic reagents
AUTHOR(S): Nishimura, Haruki; Takamatsu, Hideji
CORPORATE SOURCE: Dainippon Pharm. Co., Ltd., Osaka, Japan
SOURCE: Yakugaku Zasshi (1964), 84(9), 817-24
CODEN: YKKZAJ; ISSN: 0031-6903
DOCUMENT TYPE: Journal
LANGUAGE: Japanese
AB Treatment of DL-erythro-1-methylamino-1-phenyl-2-propanol (I) with SOCl₂ gives erythro-N-methyl-1-phenyl-2-chloropropylamine-HCl (II), m. 210° (decomposition), while treatment of I with PCl₅ gives the corresponding threo compound (IIa), m. 150-3°. Na₂S₂O₃ and II and IIa gives erythro-1-phenyl-2-methylaminopropylthiosulfonic acid (III) [m. 239° (decomposition)] and the corresponding threo compound (IIIa) [m. 199-200° (decomposition)], resp. This reaction proceeds with two Walden inversions via the intermediate aziridinium compound. Treatment of II and IIa with NaOH affords DL-threo-3-phenyl-1,2-dimethylaziridine (picrate m. 142-4°) and the corresponding erythro compound (picrate m. 97.5-8.5°), resp. Also is prepared erythro-N,N-dimethyl-1-phenyl-2-chloropropylamine-HCl (m. 169-70°) by chlorination of erythro-1-dimethylamino-1-phenyl-2-propanol-HCl with PCl₅; no threo-type Cl derivative is obtained.
IT 942-47-2, α-Toluenethiol, α-[1-(dimethylamino)ethyl]-, erythro- (preparation of)
RN 942-47-2 CAPLUS
CN α-Toluenethiol, α-[1-(dimethylamino)ethyl]-, erythro- (8CI) (CA INDEX NAME)

Relative stereochemistry.



L13 ANSWER 15 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
ACCESSION NUMBER: 1965:8904 CAPLUS
DOCUMENT NUMBER: 62:8904
ORIGINAL REFERENCE NO.: 62:1588a-c
TITLE: Phenylmercaptoalkylamines. III. Hofmann degradation of 1-phenyl-2-dimethylaminopropanethiol quaternary salts
AUTHOR(S): Nishimura, Haruki; Takamatsu, Hideji
CORPORATE SOURCE: Dainippon Pharm. Co., Ltd., Osaka, Japan
SOURCE: Yakugaku Zasshi (1964), 84(9), 811-17
CODEN: YKKZAJ; ISSN: 0031-6903
DOCUMENT TYPE: Journal
LANGUAGE: Japanese
AB Na₂S₂O₃ and L-(+)-threo-N,N-dimethyl-1-chloro-1-phenyl-2-propylamine-HCl, followed by hydrolysis, gave (+)-1-phenyl-2-dimethylaminopropanethiol (I), which was then converted into the methiodide and treated with NaOH to form (+)-1,2-epithiopropylbenzene (II), b₁₀ 100°, which was polymerized to give a polymer, m. 255-6°. Treatment of D-(+)-erythro-1,2-epoxypropylbenzene with KSCN gave L-(-)-erythro-1,2-epithiopropylbenzene, b₇ 92-3°, [α]_{20D}-21.4° (c 2.21, MeOH), which was found to be the antipode of II. II belongs to the D-(+)-erythro series and I,


to the L-(+)-threo series. The (-)-amino thiol, similarly derived from L-(-)-erythro-N,N-dimethyl-1-chloro-1-phenyl-2-propylamine-HCl, was found to belong to the L-(-)-erythro series and that D-(+)-threo-1,2-epithiopropylbenzene (III) is derived from it. The steric configuration of II and III was also determined from their N.M.R. spectra. Hofmann degradation of the quaternary salt of 1-phenyl-2-dimethylaminoethanethiol also gave the same result. II and III underwent desulfurization by heating to give trans- β -methylstyrene.

IT 942-50-7, α -Toluenethiol, α -[1-(dimethylamino)ethyl]-, hydrochloride, L-(+)-threo- (preparation of)

RN 942-50-7 CAPLUS

CN Benzenemethanethiol, α -[1-(dimethylamino)ethyl]-, hydrochloride, [S-(R*,R*)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl

L13 ANSWER 16 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1965:8903 CAPLUS

DOCUMENT NUMBER: 62:8903

ORIGINAL REFERENCE NO.: 62:1587h,1588a

TITLE: Phenylmercaptoalkylamines. II. Configuration of 1-phenyl-2-alkylaminopropanethiol

AUTHOR(S): Nishimura, Haruki; Takamatsu, Hideji

CORPORATE SOURCE: Dainippon Pharm. Co., Ltd., Osaka, Japan

SOURCE: Yakugaku Zasshi (1964), 84(9), 806-11

CODEN: YKKZAJ; ISSN: 0031-6903

DOCUMENT TYPE: Journal


LANGUAGE: Japanese

AB From the strength of the H bond between N and SH in the ir absorption spectra of (+)- (I) and (-)-1-phenyl-2-dimethylaminopropanethiol (Ia), obtained by treating Na₂S₂O₃ with L-(+)-threo- (II) and L-(-)-erythro-N,N-dimethyl-1-chloro-1-phenyl-2-propylamine-HCl (IIa) followed by hydrolysis, it was assumed that I belongs to the threo series and Ia, to the erythro series. I is also obtained by treating AcSK or NaSH with II or by treating (NH₂)₂CS with II followed by alkali treatment of the isothiuronium salt formed. Similarly, the steric configurations of 2 1-phenyl-2-methylaminopropanethiols and 2-imino-3,4-dimethyl-5-phenylthiazolidines, derived from L-(+)-threo-(III) and L-(-)-erythro-N-methyl-1-chloro-1-phenyl-2-propylamine hydrochlorides (IIIa) by use of foregoing reagents, were determined. Use of nucleophilic reagents to II, IIa, III, and IIIa afforded derivs. without change in their steric configuration.

IT 942-48-3, α -Toluenethiol, α -[1-(dimethylamino)ethyl]-, L-(+)-threo- 942-49-4, α -Toluenethiol, α -[1-(dimethylamino)ethyl]-, L-(-)-erythro- 942-50-7,

LANGUAGE: Unavailable
 PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	US 3131189		19640428	US	19611016 <--
GI	For diagram(s), see printed CA Issue.				
AB	Carbyl sulfate (I), prepared by the reaction of 2 moles SO ₃ and 1 mole ethylene, reacted with a tertiary amine to form betaines. Thus 1.5 g. pyridine (II) in 10 ml. ethylene dichloride was added to 3 g. I in 30 ml. ethylene dichloride (the reaction was exothermic), the liquid decanted from the precipitate, and the precipitate covered with petr. ether and cooled to give				
	IIa (R = R ₁ = H), m. 250-5° (HCONMe ₂). I was also treated with the following to form betaines: quinoline, acridine, trimethylamine, and dimethylaniline (III). Also reported without details were: IIa (R = Ph, R ₁ = H); Et ₃ NCH ₂ CH ₂ SO ₃ ; IIa (R = R ₁ = Me); and PhNMe ₂ CMe ₂ CMe ₂ SO ₃ . These compds. are useful intermediates for the preparation of detergents. (Cf. U.S. 2,666,788, or Brit. 686,061.)				
IT	97176-62-0, Ammonium, dimethylphenyl(1,1,2-trimethyl-2-sulfopropyl), hydroxide, inner salt (preparation of)				
RN	97176-62-0 CAPLUS				
CN	Dimethylphenyl(1,1,2-trimethyl-2-sulfopropyl)ammonium hydroxide, inner salt (7CI) (CA INDEX NAME)				



=> d l14 ibib abs hitstr tot

L14 ANSWER 1 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 2001:821889 CAPLUS
 DOCUMENT NUMBER: 136:118295
 TITLE: Stereoselective synthesis of δ -lactones from 5-oxoalkanals via one-pot sequential acetalization, Tishchenko reaction, and lactonization by cooperative catalysis of samarium ion and mercaptan
 AUTHOR(S): Hsu, Jue-Liang; Fang, Jim-Min
 CORPORATE SOURCE: Department of Chemistry, National Taiwan University, Taipei, 106, Taiwan
 SOURCE: Journal of Organic Chemistry (2001), 66(25), 8573-8584
 CODEN: JOCEAH; ISSN: 0022-3263
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB By the synergistic catalysis of samarium ion and mercaptan, a series of 5-oxoalkanals was converted to (substituted) δ -lactones in efficient and stereoselective manners. This one-pot procedure comprises a sequence

of acetalization, Tishchenko reaction and lactonization. The deliberative use of mercaptan, by comparison with alc., is advantageous to facilitate the catalytic cycle. The reaction mechanism and stereochem. are proposed and supported by some exptl. evidence. Such samarium ion/mercaptan cocatalyzed reactions show the feature of remote control, which is applicable to the asym. synthesis of optically active δ -lactones. This study also demonstrates the synthesis of two insect pheromones, (2S,5R)-2-methylhexanolide and (R)-hexadecanolide, as examples of a new protocol for asym. reduction of long-chain aliphatic ketones.


IT 166031-49-8P

RL: SPN (Synthetic preparation); PREP (Preparation)
(one-pot sequential acetalization, Tishchenko reaction, and lactonization by the promotion of samarium ion and mercaptans in stereoselective synthesis of δ -lactones from 5-oxoalkanal)

RN 166031-49-8 CAPLUS

CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
(α R, β S) - (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 104 THERE ARE 104 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L14 ANSWER 2 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2001:508661 CAPLUS

DOCUMENT NUMBER: 135:256816

TITLE: A purely synthetic, diversity amenable version of norephedrine thiols for the highly enantioselective diethylzinc addition to aldehydes

AUTHOR(S): Jimeno, Ciril; Moyano, Albert; Pericas, Miquel A.; Riera, Antoni

CORPORATE SOURCE: Unitat Recerca Sintesi Asimetrica, Dep. Quim. Org., Universitat de Barcelona, Barcelona, E-08028, Spain

SOURCE: Synlett (2001), (7), 1155-1157
CODEN: SYNLES; ISSN: 0936-5214

PUBLISHER: Georg Thieme Verlag

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 135:256816

AB A new β -amino thiol arising from purely synthetic yet enantiopure amino alcs. has been prepared and successfully used in the addition of diethylzinc to aromatic aldehydes, yielding secondary alcs. in ee's up to 99%.

IT 361543-74-0P

RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation); USES (Uses)


(enantioselective diethylzinc addition to aldehydes catalyzed by β -amino thiols)

RN 361543-74-0 CAPLUS

03/14/2005 10807710.trn

CN 1-Piperidineethanethiol, α -phenyl- β -[(triphenylmethoxy)methyl]-
, (α R, β S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 3 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1999:296019 CAPLUS

DOCUMENT NUMBER: 130:312007

TITLE: A concise synthesis of unnatural (+)-5-epi-nojirimycin- δ -lactam via asymmetric reduction of a meso-imide

AUTHOR(S): Kang, Jahyo; Lee, Choon Woo; Lim, Geun Jho; Cho, Byung Tae

CORPORATE SOURCE: Department of Chemistry, Sogang University, Seoul, 121-742, S. Korea

SOURCE: Tetrahedron: Asymmetry (1999), 10(4), 657-660

CODEN: TASYE3; ISSN: 0957-4166

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 130:312007

AB Nojirimycin- δ -lactam skeleton was synthesized by asym. reduction of a cyclic triacetyloxy meso imide with a chiral β -amino thiol ligand. The resulting product was converted to unnatural (+)-5-epi-nojirimycin- δ -lactam.

IT 166031-49-8


RL: RCT (Reactant); RACT (Reactant or reagent)

(a concise synthesis of unnatural (+)-epi-nojirimycin- δ -lactam via asym. reduction of a meso-imide)

RN 166031-49-8 CAPLUS

CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
(α R, β S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 4 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1998:801743 CAPLUS
 DOCUMENT NUMBER: 130:153758
 TITLE: Asymmetric synthesis of (diene)Fe(CO)₃ complexes by a catalytic enantioselective alkylation using dialkylzincs
 AUTHOR(S): Takemoto, Yoshiji; Baba, Yasutaka; Honda, Asami; Nakao, Syusuke; Noguchi, Izumi; Iwata, Chuzo; Tanaka, Tetsuaki; Ibuka, Toshiro
 CORPORATE SOURCE: Graduate School of Pharmaceutical Sciences, Kyoto University, Kyoto, 606-8501, Japan
 SOURCE: Tetrahedron (1998), 54(51), 15567-15580
 CODEN: TETRAB; ISSN: 0040-4020
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 130:153758

AB The reaction of meso-(2,4-hexadien-1,6-dial)Fe(CO)₃ complex (1) with several alkylzincs in the presence of 50 mol% of (S)-(+)-diphenyl(1-methylpyrrolidin-2-yl)methanol proceeded with high enantiotopic group- and diastereotopic face-selectivity to give (2R,6S)-alc. complexes as major products, except in the case with dimethylzinc (>90% de and >98% ee). On the other hand, the methylation of 1 with Me₂Zn proceeded with high enantioselectivity by adding 1.8 equivalent of Ti(Oi-Pr)₄ in the presence of 3 mol% of (S,S)-1,2-bis(trifluoromethylsulfonamide)cyclohexane (82% de, 96% ee). The enantioselective alkylation was also applied to the kinetic resolution of racemic (sorbic aldehyde)Fe(CO)₃ complex.


IT 166031-49-8

RL: CAT (Catalyst use); USES (Uses)
 (asym. synthesis of diene iron tricarbonyl complexes by catalytic enantioselective alkylation using dialkylzincs)

RN 166031-49-8 CAPLUS

CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
 (α R, β S) - (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 5 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1998:632164 CAPLUS
 DOCUMENT NUMBER: 129:343594
 TITLE: Structural chemistry of methyl- and allylpalladium(II) complexes containing chiral thioether auxiliaries
 AUTHOR(S): Boog-Wick, Karin; Pregosin, Paul S.; Woerle, Michael; Albinati, Alberto
 CORPORATE SOURCE: Lab. Anorganische Chem., ETH Zentrum Zuerich, Zurich,


CH-8092, Switz.
 SOURCE: Helvetica Chimica Acta (1998), 81(9),
 1622-1633
 CODEN: HCACAV; ISSN: 0018-019X
 PUBLISHER: Verlag Helvetica Chimica Acta AG
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 129:343594

AB The synthesis and mol. structures of two [PdCl(Me)] complexes each containing a different chiral N,S-chelate based on {[dihydrooxazolyl]phenyl}methyl}thiogluco-
 glucose backbones, i.e., chloro({2-[(4S)-4,5-dihydro-4-isopropoxyloxazol-
 2-yl-κN]phenyl}methyl 2,3,4,6-tetra-O-acetyl-1-(thio-κS)-
 β-D-glucopyranoside)methylpalladium(II) and a [Pd(η³-C₃H₅)(PS)]⁺
 cation in which the P,S-chelate stems from a phosphinoferrocene and
 thioephedrine-derived thioether donor as well as [(S)-1-(diphenylphosphino-
 κP)-2-((1R)-1-[[[(1R,2S)-1-phenyl-2-(piperidin-1-yl)propyl]thio-
 κS}ethyl)ferrocene](η³-prop-2-enyl)palladium
 trifluoromethanesulfonate are reported. In the methylpalladium compds.
 the thiogluco-κS moiety is pseudo-axial, whereas in the allyl
 complex, the thioephedrine-κS moiety is markedly pseudo-equatorial.
 It is suggested, based on these results, that the shape (chiral pocket) of
 such coordinated chiral thioethers may not be readily predictable.

IT 166031-49-8
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of methyl- and allylpalladium complexes with chiral thioether
 moieties)

RN 166031-49-8 CAPLUS
 CN 1-Piperidineethanethiol, β-methyl-α-phenyl-,
 (αR,βS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



L14 ANSWER 6 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1998:482774 CAPLUS
 DOCUMENT NUMBER: 129:216335
 TITLE: Asymmetric synthesis of a new cylindrically chiral and
 air-stable ferrocenyldiphosphine and its application
 to rhodium-catalyzed asymmetric hydrogenation
 AUTHOR(S): Kang, Jahyo; Lee, Jun Hee; Ahn, Sung Hoon; Choi, Jung
 Sun
 CORPORATE SOURCE: Department of Chemistry and Organic Chemistry Research
 Center, Sogang University, Seoul, 121-742, S. Korea
 SOURCE: Tetrahedron Letters (1998), 39(31),
 5523-5526
 CODEN: TELEAY; ISSN: 0040-4039
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 129:216335

AB The novel, cylindrically chiral air-stable (S,S)-1,1'-bis(diphenylphosphino)-2,2'-di-3-pentylferrocene [(S,S)-FerroPHOS] ligand was prepared and its in situ rhodium complexes have been applied to asym. hydrogenation. High reactivity and selectivity have been realized in hydrogenation of various dehydroamino acid derivs.


IT 166031-49-8

RL: RCT (Reactant); RACT (Reactant or reagent)
(asym. synthesis of a new cylindrically chiral and air stable ferrocenyldiphosphine and its application to rhodium catalyzed asym. hydrogenation)

RN 166031-49-8 CAPLUS

CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
(α R, β S) - (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 7 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1998:362994 CAPLUS

DOCUMENT NUMBER: 129:122430

TITLE: Examination of bidentate thiol derivatives as ligands in the Ni-catalyzed asymmetric conjugate addition of diethylzinc to enones

AUTHOR(S): Kang, Jahyo; Kim, Joo In; Lee, Jae Hoon; Kim, Hyo - Jung; Byun, Yong Hun

CORPORATE SOURCE: Department of Chemistry, Sogang University, Seoul, 121-742, S. Korea

SOURCE: Bulletin of the Korean Chemical Society (1998), 19(5), 601-603

CODEN: BKCSDE; ISSN: 0253-2964


PUBLISHER: Korean Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 129:122430


GI



AB Asym. conjugate addition of diethylzinc to (E)-PhCH:CHCOPh was catalyzed by Ni(II) complexes, e.g. Ni(acac)₂ or NiCl₂, in presence of bidentate thiol

ligands, e.g. I. The major enantiomer obtained was (R)-PhCH₂EtCH₂COPh.
 IT 166031-49-8
 RL: CAT (Catalyst use); USES (Uses)
 (asym. conjugate addition of diethylzinc to enones catalyzed by Ni(II) and
 bidentate thiol ligands)
 RN 166031-49-8 CAPLUS
 CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
 (α R, β S) - (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




REFERENCE COUNT: 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 8 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1998:272187 CAPLUS
 DOCUMENT NUMBER: 129:54427
 TITLE: The interaction of chiral amino thiols with organozinc
 reagents and aldehydes: a mechanism of amino
 thiol-catalyzed addition of organozinc reagents to
 aldehydes
 AUTHOR(S): Kang, Jahyo; Kim, Jin Bum; Kim, Jeeyoung; Lee,
 Duckhwan
 CORPORATE SOURCE: Department of Chemistry, Sogang University, Seoul,
 121-742, S. Korea
 SOURCE: Bulletin of the Korean Chemical Society (1998
), 19(4), 475-481
 CODEN: BKCSDE; ISSN: 0253-2964
 PUBLISHER: Korean Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Details of various equilibrium involved in the reactions of oxaza- and
 thiazazincolidine catalysts, generated from either β -amino alc. or
 β -amino thiol, with benzaldehyde were studied by colligative
 measurements. The coordination of diethylzinc prior to the coordination
 of aldehyde is essential for high enantioselectivity of the thiol
 catalyzed reaction. The probable origin of asym. nonlinearity is also
 presented.


IT 166031-49-8 188711-05-9
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (mechanism of amino thiol-catalyzed addition of organozinc reagents to
 aldehydes)
 RN 166031-49-8 CAPLUS
 CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
 (α R, β S) - (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 188711-05-9 CAPLUS
 CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
 (α R, β S)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 9 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1997:213617 CAPLUS
 DOCUMENT NUMBER: 126:277632
 TITLE: Enantioselective catalytic reduction of dihydroisoquinoline derivatives
 AUTHOR(S): Kang, Jahyo; Kim, Jin Bum; Cho, Kwi Hyung; Cho, Byung Tae
 CORPORATE SOURCE: Department of Chemistry, Sogang University, Seoul, 121-742, S. Korea
 SOURCE: Tetrahedron: Asymmetry (1997), 8(5), 657-660
 CODEN: TASYE3; ISSN: 0957-4166
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 126:277632
 GI




AB Zinc complexes I [RR = (CH₂)₅, R = Me] were shown to be excellent catalysts for enantioselective reduction of dihydroisoquinolines II (R₁ = Me, 3,4-dimethoxy-, 3,4,5-trimethoxybenzyl, 3,4-dimethoxyphenyl) with

BH3·THF to the corresponding (R)-tetrahydroisoquinolines III with good enantioselectivity.


IT 166031-49-8
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (enantioselective catalytic reduction of dihydroisoquinoline derivs.)
 RN 166031-49-8 CAPLUS
 CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
 (α R, β S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L14 ANSWER 10 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1997:133573 CAPLUS
 DOCUMENT NUMBER: 126:263720
 TITLE: The effects of sulfur substitution in chiral amino thiols on the enantioselective addition of organozinc reagents to aldehydes: a novel method for estimation of free energies of dimerization in monomer-dimer equilibria
 AUTHOR(S): Kang, Jahyo; Kim, Jin Bum; Kim, Jeong Whan; Lee, Duckhwan
 CORPORATE SOURCE: Dep. of Chemistry, Sogang University, Seoul, 121-742, S. Korea
 SOURCE: Journal of the Chemical Society, Perkin Transactions 2: Physical Organic Chemistry (1997), (2), 189-194
 CODEN: JCPKBH; ISSN: 0300-9580
 PUBLISHER: Royal Society of Chemistry
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



I

AB Differences between the thiol ligand I (X = S) and the corresponding alc. ligand (X = O) were observed in the catalytic asym. alkylation of benzaldehyde with diethylzinc. The thiol ligand was superior for reaction rate, enantioselectivity and asym. amplification. The effects of chiral amino thiols are discussed and compared with the results of chiral amino

alc. counterparts. The quant. and thermodyn. aspects of the monomer-dimer equilibrium involved in thiazazincolidine or oxazazincolidine catalysts have also been studied on the basis of colligative properties.

IT 166031-49-8 188711-05-9 188711-34-4


RL: CAT (Catalyst use); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); PROC (Process); RACT (Reactant or reagent); USES (Uses)

(effect of sulfur substitution in chiral amino thiols on enantioselective addition of organozinc reagents to aldehydes, and colligative estimation of free energies of dimerization in monomer-dimer equilibrium)

RN 166031-49-8 CAPLUS

CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
(α R, β S) - (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 188711-05-9 CAPLUS

CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
(α R, β S)-rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 188711-34-4 CAPLUS

CN 1-Piperidineethanethiol, β -methyl- α -phenyl-, [S-(R*,S*)] - (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:

16

THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L14 ANSWER 11 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1997:39785 CAPLUS
 DOCUMENT NUMBER: 126:131036
 TITLE: Chiral β -amino thiol catalysts for the
 enantioselective addition of diethylzinc to aldehydes
 AUTHOR(S): Kang, Jahyo; Kim, Jeong Whan; Lee, Jun Won; Kim, Dong
 Soo; Kim, Joo In
 CORPORATE SOURCE: Dep. Chem., Sogang Univ., Seoul, 121-742, S. Korea
 SOURCE: Bulletin of the Korean Chemical Society (1996
), 17(12), 1135-1142
 CODEN: BKCSDE; ISSN: 0253-2964
 PUBLISHER: Korean Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Reaction of diethylzinc with α -branched aldehydes in the presence of
 a catalytic amount (5 mol %) of various β -amino thiols in toluene or
 ether provided the corresponding secondary alcs. in outstanding ee.
 Detailed preparative procedure for the β -amino thiols are presented.


IT 160011-80-3P
 RL: CAT (Catalyst use); RCT (Reactant); SPN (Synthetic preparation); PREP
 (Preparation); RACT (Reactant or reagent); USES (Uses)
 (enantioselective addition of diethylzinc to aldehydes using chiral
 β -amino thiol catalysts)
 RN 160011-80-3 CAPLUS
 CN 1-Piperidineethanethiol, α,β -diphenyl-, ($\alpha R,\beta S$)-
 (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



IT 166031-49-8P
 RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);
 USES (Uses)
 (enantioselective addition of diethylzinc to aldehydes using chiral
 β -amino thiol catalysts)
 RN 166031-49-8 CAPLUS
 CN 1-Piperidineethanethiol, β -methyl- α -phenyl-,
 ($\alpha R,\beta S$)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L14 ANSWER 12 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1995:194984 CAPLUS
 DOCUMENT NUMBER: 122:55341
 TITLE: Enantioselective addition of diethylzinc to aldehydes
 catalyzed by a drug-unrelated chiral amino thiol and
 the corresponding disulfide
 AUTHOR(S): Kang, Jahyo; Kim, Dong Soo; Kim, Joo In
 CORPORATE SOURCE: Department Chemistry, Sogang University, Seoul,
 121-742, S. Korea
 SOURCE: Synlett (1994), (10), 842-4
 CODEN: SYNLES; ISSN: 0936-5214
 PUBLISHER: Thieme
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 122:55341
 GI




AB Reaction of diethylzinc with aldehydes in the presence of a catalytic amount of a β -amino thiol I (5 mol %) and the disulfide II (2.5 mol %) in toluene at 0° provided the corresponding secondary alcs. in excellent ee's.

IT- 160011-80-3P-
 RL: CAT (Catalyst use); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (enantioselective addition of diethylzinc with aldehydes catalyzed by chiral amino thiol and disulfide)

RN 160011-80-3 CAPLUS

CN 1-Piperidineethanethiol, α,β -diphenyl-, (α R, β S)-
 (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).




L14 ANSWER 13 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1995:86424 CAPLUS
 DOCUMENT NUMBER: 123:142957
 TITLE: Enantioselective addition of diethylzinc to

AUTHOR(S): Kang, Jahyo; Lee, Jun Won; Kim, Joo In
 CORPORATE SOURCE: Department of Chemistry, Sogang University, Seoul, 121-742, S. Korea
 SOURCE: Journal of the Chemical Society, Chemical Communications (1994), (17), 2009-10
 CODEN: JCCCAT; ISSN: 0022-4936
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 123:142957
 AB Reaction of diethylzinc with α -branched aldehydes in the presence of a catalytic amount of (1R,2S)-(-)-1-phenyl-2-piperidinopropane-1-thiol provided the corresponding secondary alcs. in almost 100% enantiomeric excess.
 IT 166031-49-8P
 RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation); USES (Uses)
 (best catalyst; as ligand catalyst for enantioselective addition of diethylzinc to α -branched aldehydes)
 RN 166031-49-8 CAPLUS
 CN 1-Piperidineethanethiol, β -methyl- α -phenyl-, (α R, β S) - (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



L14 ANSWER 14 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1993:560233 CAPLUS
 DOCUMENT NUMBER: 119:160233
 TITLE: Reactions of carbanions generated from 2-(dialkylamino)-2-phenylacetonitriles with disubstituted acetylenes
 AUTHOR(S): Zdrojewski, Tadeusz; Jonczyk, Andrzej
 CORPORATE SOURCE: Dep. Chem., Tech. Univ. (Politechnika), Warsaw, 00-662, Pol.
 SOURCE: Liebigs Annalen der Chemie (1993), (4), 375-8
 CODEN: LACHDL; ISSN: 0170-2041
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



II

AB Reactions of aminonitriles $\text{PhCH}(\text{NR}_{12})\text{CN}$ [$\text{R}_{12} = (\text{CH}_2)_5, (\text{CH}_2)_4, (\text{CH}_2)_2\text{O}(\text{CH}_2)_2, (\text{CH}_2)_2\text{S}(\text{CH}_2)_2, (\text{CH}_2)_2\text{NMe}(\text{CH}_2)_2$] with MeC.tplbond.CR_2 ($\text{R}_2 = \text{SMe}, \text{Ph}$), carried out in DMSO in the presence of powdered sodium hydroxide and benzyltriethylammonium chloride (TEBAC) as a catalyst, afford either $\text{PhC}(\text{NR}_{12})(\text{CN})\text{CMe:CHR}_2$ (I) as the E/Z isomer mixture or the pure E isomer or a mixture of I ($\text{R}_2 = \text{SMe}$) and dipiperidinopentenitrile II; unmasking of the carbonyl group in I [$\text{R}_{12} = (\text{CH}_2)_5, \text{R}_2 = \text{Ph}$] and II gives ketones PhCOCMe:CHPh and $\text{PhCOCHMeCH}(\text{SMe})\text{COPh}$, resp.


IT 150179-13-8P 150179-14-9P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and cleavage of)

RN 150179-13-8 CAPLUS

CN 1-Piperidineacetonitrile, α -[2-methyl-1-(methylthio)-3-phenyl-3-(1-piperidinyl)-2-propenyl]- α -phenyl-, (R^*, R^*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.
Double bond geometry unknown.



RN 150179-14-9 CAPLUS

CN 1-Piperidineacetonitrile, α -[2-methyl-1-(methylthio)-3-phenyl-3-(1-piperidinyl)-2-propenyl]- α -phenyl-, (R^*, S^*)- (9CI) (CA INDEX NAME)

Relative stereochemistry.
Double bond geometry unknown.



L14 ANSWER 15 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1991:228476 CAPLUS

DOCUMENT NUMBER: 114:228476

TITLE: Synthesis and study of derivatives of
3-(alkylamino)-2-(methylthio)carboxylic acids
Vidugiriene, V.; Valaviciene, J.; Rasteikiene, L.
CORPORATE SOURCE: Inst. Biokhim., Vilnius, USSR
SOURCE: Chemija (1990), (2), 101-6
CODEN: CHMJES; ISSN: 0235-7216

DOCUMENT TYPE: Journal

LANGUAGE: Russian

OTHER SOURCE(S): CASREACT 114:228476

AB Treating $\text{Z-RCH:C}(\text{SMe})\text{COX}$ ($\text{R} = \text{Ph}, \text{Me}, \text{X} = \text{NHPh}$; $\text{R} = \text{Ph}, \text{X} = \text{OMe}$) and

RCHClCH(SMe)COX (same R, X) with 2-4 equiv HA (A = piperidino, morpholino, cyclohexylamino, ethylenimino, NHC5H11-n) gave 10 corresponding RCHACH(SMe)COX (I) as erythro-threo mixts. I had low-to-moderate toxicity and antitumor activity, with I (R = Ph, A = piperidino, X = NHPh) showing the best profile.

IT 133508-72-2P 133508-73-3P 133508-80-2P


133508-81-3P 133612-17-6P 133612-23-4P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation, toxicity, and antineoplastic activity of)

RN 133508-72-2 CAPLUS

CN 1-Piperidinepropanoic acid, α -(methylthio)- β -phenyl-, methyl ester, (R*,R*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 133508-73-3 CAPLUS

CN 1-Piperidinepropanoic acid, α -(methylthio)- β -phenyl-, methyl ester, (R*,S*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 133508-80-2 CAPLUS

CN 1-Piperidinepropanamide, α -(methylthio)-N, β -diphenyl-, (R*,R*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 133508-81-3 CAPLUS

CN 1-Piperidinepropanamide, α-(methylthio)-N,β-diphenyl-, (R*,S*)-
(9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 133612-17-6 CAPLUS

CN 1-Piperidinepropanoic acid, β-methyl-α-(methylthio)-, methyl
ester, (R*,R*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.




RN 133612-23-4 CAPLUS

CN 1-Piperidinepropanoic acid, β-methyl-α-(methylthio)-, methyl
ester, (R*,S*)- (9CI) (CA INDEX NAME)

Relative stereochemistry.



L14 ANSWER 16 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
 ACCESSION NUMBER: 1991:206268 CAPLUS
 DOCUMENT NUMBER: 114:206268
 TITLE: Nucleophilic addition of amines to derivatives of unsaturated acids containing 2-alkyl(phenyl)thio groups
 AUTHOR(S): Talaikyte, Z.; Vidugiriene, V.; Rasteikiene, L.
 CORPORATE SOURCE: Inst. Biokhim., Vilnius, USSR
 SOURCE: Chemija (1990), (2), 93-100
 CODEN: CHMJES; ISSN: 0235-7216
 DOCUMENT TYPE: Journal
 LANGUAGE: Russian
 GI



I


AB The title reaction of piperidine or morpholine with $\text{RCH}:\text{C}(\text{SR}_1)\text{COX}$ ($\text{R} = \text{Me}$, Ph ; $\text{R}_1 = \text{Me}$, Ph , $\text{CH}_2\text{CH}_2\text{Cl}$; $\text{X} = \text{NHPh}$, OMe) is nonstereospecific and gives mixts. of derivs. of erythro- and threo-butanoic acid I ($\text{R} = \text{Me}$, $\text{R}_2 = \text{Me}$, piperidino- or morpholinoethyl; $\text{Y} = \text{CH}_2$, O) or -phenylpropanoic acid I ($\text{R} = \text{Ph}$).

IT 133508-66-4P 133508-67-5P 133508-72-2P
 133508-73-3P 133508-80-2P 133508-81-3P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)

RN 133508-66-4 CAPLUS

CN 1-Piperidinepropanamide, β -methyl- α -(methylthio)-N-phenyl-,
 (R^* , R^*) - (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 133508-67-5 CAPLUS

CN 1-Piperidinepropanamide, β -methyl- α -(methylthio)-N-phenyl-,
(R*,S*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 133508-72-2 CAPLUS

CN 1-Piperidinepropanoic acid, α -(methylthio)- β -phenyl-, methyl
ester, (R*,R*)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 133508-73-3 CAPLUS


CN 1-Piperidinepropanoic acid, α -(methylthio)- β -phenyl-, methyl
ester, (R*,S*)- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 133508-80-2 CAPLUS
CN 1-Piperidinepropanamide, α -(methylthio)-N, β -diphenyl-, (R*,R*)-
(9CI) (CA INDEX NAME)

Relative stereochemistry.



RN 133508-81-3 CAPLUS
CN 1-Piperidinepropanamide, α -(methylthio)-N, β -diphenyl-, (R*,S*)-
(9CI) (CA INDEX NAME)


Relative stereochemistry.



L14 ANSWER 17 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN
ACCESSION NUMBER: 1990:458617 CAPLUS
DOCUMENT NUMBER: 113:58617
TITLE: Reactions of organic anions. 168. Reactions of
2-(dialkylamino)arylacetonitriles with acetylenes
under basic conditions. A simple synthesis of
substituted mono- and diketones
AUTHOR(S): Zdrojewski, T.; Jonczyk, A.
CORPORATE SOURCE: Dep. Chem., Tech. Univ., Warsaw, PL-00-662, Pol.

03/14/2005 10807710.trn

SOURCE: Synthesis (1990), (3), 224-33
CODEN: SYNTBF; ISSN: 0039-7881
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 113:58617
AB The reaction of $\text{RCH}(\text{CN})\text{NR}_{12}$ (I; R = Ph, 4-MeC₆H₄, 4-MeOC₆H₄; R₁ = Me; NR₁₂ = piperidino, morpholino, etc.) with $\text{R}_2\text{C}=\text{CH}\text{CH}_2$ (II; R₂ = Ph, MeS) gave $\text{R}_1\text{NCR}(\text{CN})\text{CH}=\text{CHR}_2$ (III) and/or $\text{R}_1\text{NCR}(\text{CN})\text{CHR}_2\text{CH}=\text{CHR}_2$; the product depended on the basicity of the amino group in III. I also added to C-1 of II (R₂ = EtO) to give $\text{R}_1\text{NCR}(\text{CN})\text{C}(\text{OEt})\text{CH}_2$. All these products could be hydrolyzed to give mono- or diketones.
IT 128407-40-9P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and hydrolysis of)
RN 128407-40-9 CAPLUS
CN 1-Piperidineacetonitrile, α -[1-(methylthio)-3-phenyl-3-(1-piperidinyl)-2-propenyl]- α -phenyl- (9CI) (CA INDEX NAME)



=> log y

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
173.54	677.96

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-24.82	-27.01

CA SUBSCRIBER PRICE

STN INTERNATIONAL LOGOFF AT 14:27:52 ON 14 MAR 2005